

Section 2.3 Q4: Consider the diffusion eq. $u_t = u_{xx}$ in $(x, t) \in (0, 1) \times (0, \infty)$ with $u(0, t) = u(1, t) = 0$ and $u(x, 0) = 4x(1-x)$.

a) Show that $0 < u(x, t) < 1 \quad \forall t > 0, 0 < x < 1$.

b) Show that $u(x, t) = u(1-x, t) \quad \forall t \geq 0, 0 \leq x \leq 1$.

c) Use the energy method to show that $\int_0^1 u(x, t)^2 dx$ is a strictly decreasing function of t .

Denote $R = [0, 1] \times [0, \infty)$,

$\Gamma = \{\text{bottom}\} \cup \{\text{left side}\} \cup \{\text{right side}\}$

a) By the strong maximum principle

the max of $u(x, t)$ in R

is achieved on the boundary Γ .

On the sides $u=0$. On the

bottom $u(x, 0) = 4x(1-x)$,

$$u\left(\frac{1}{2}, 0\right) = 4 \cdot \frac{1}{2} \cdot \frac{1}{2} = 1.$$

It is easy to see that this is the max.

So $u(x, t) < 1$ (strict $<$) inside R , i.e. for


$$x \in (0, 1), t > 0.$$

Similarly, by the strong minimum principle, u achieves its

minimum on Γ . Since $u \geq 0$ on Γ , it must hold that

$u(x, t) > 0$ (strict $>$) inside R . So inside R

$$0 < u(x, t) < 1.$$



b) Let $v(x,t) = u(1-x,t)$. Then:

$$v_t = u_t, \quad v_x = -u_x, \quad v_{xx} = -(-u_x)_x = u_{xx}.$$

Hence $v_t - v_{xx} = u_t - u_{xx} = 0$.

Moreover: $v(0,t) = u(1,t) = 0$

$$v(1,t) = u(0,t) = 0$$

$$v(x,0) = u(1-x,0) = 4(1-x)x$$

So v solves the same problem like u . We know that solutions are unique ("Uniqueness of Solutions" theorem) so that u and v must be the same: $u(x,t) = v(x,t) = u(1-x,t)$ for all $t \geq 0$ and $0 \leq x \leq 1$.

c) The **energy method** is the method where we multiply the eq. by u and integrate:

The eq. is: $u_t - u_{xx} = 0$

Multiply: $u(u_t - u_{xx}) = 0$

Integrate: $\int_0^1 u(x,t) [u_t(x,t) - u_{xx}(x,t)] dx = 0$.

$$\begin{aligned} \Rightarrow 0 &= \int_0^1 u u_t dx - \int_0^1 u u_{xx} dx \xrightarrow{\text{int. by parts}} \\ &= \frac{1}{2} \frac{d}{dt} \int_0^1 u^2 dx + \int_0^1 u_x^2 dx - \underbrace{[u u_x]_{x=0}^1}_{=0 \text{ since } u(0,t)=u(1,t)=0} \end{aligned}$$

$$\Rightarrow \frac{d}{dt} \int_0^1 u^2 dx = -2 \int_0^1 u_x^2 dx \leq 0$$

But we know that $\int_0^1 u_x^2 dx > 0$ strictly (i.e. it is not 0).

How do we know this? From part (a) we know that $0 < u < 1$ inside \mathcal{R} , yet $u=0$ on the sides.

This means that it is impossible for u_x to always be 0 along lines of constant t .

Hence $\int_0^l u^2 dx$ strictly decreases in time.

Section 2.3 Q6: Prove the comparison principle for the diffusion eq: if u and v are two solutions and if $u \leq v$ for $t=0$, $x=0$, $x=l$, then $u \leq v$ for $t \geq 0$ and $x \in [0, l]$.

Define $w = u - v$.


Then $w \leq 0$ on

$$\Gamma = \{\text{bottom}\} \cup \{\text{right}\} \cup \{\text{left}\}.$$

By linearity of the diffusion eq, w is also a solution.

By the maximum principle, $w \leq 0$ within the infinite rectangle $R = [0, l] \times [0, \infty)$.

$$\text{So } u - v = w \leq 0 \quad \longrightarrow \quad u \leq v \quad \text{in } R.$$



Section 2.4 Q1: Solve the diffusion eq. with the initial condition

$$\phi(x) = \begin{cases} 1 & |x| < l \\ 0 & |x| > l \end{cases}$$

We know that the formula is

$$u(x,t) = \frac{1}{\sqrt{4\pi kt}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4kt}} \phi(y) dy.$$

In our case this simplifies to

$$u(x,t) = \frac{1}{\sqrt{4\pi kt}} \int_{-l}^l e^{-\frac{(x-y)^2}{4kt}} dy$$

To express in terms of the error function, make the change of variables $p = \frac{y-x}{\sqrt{4kt}}$ so that

$$dp = \frac{dy}{\sqrt{4kt}} \rightarrow dy = \sqrt{4kt} dp$$

$$\begin{aligned} \rightarrow u(x,t) &= \frac{1}{\sqrt{\pi}} \int_{-\frac{l-x}{\sqrt{4kt}}}^{\frac{l-x}{\sqrt{4kt}}} e^{-p^2} dp \\ &= \frac{1}{\sqrt{\pi}} \int_0^{\frac{l-x}{\sqrt{4kt}}} e^{-p^2} dp - \frac{1}{\sqrt{\pi}} \int_0^{-\frac{l-x}{\sqrt{4kt}}} e^{-p^2} dp \\ &= \frac{1}{2} \operatorname{Erf}\left(\frac{l-x}{\sqrt{4kt}}\right) - \frac{1}{2} \operatorname{Erf}\left(\frac{-l-x}{\sqrt{4kt}}\right). \end{aligned}$$

Section 2.4 Q6: Compute $\int_0^{\infty} e^{-x^2} dx$.

We've done this in class!

Section 2.4 Q7: Show that $\int_{-\infty}^{\infty} e^{-p^2} dp = \sqrt{\pi}$ and that $\int_{-\infty}^{\infty} \zeta(x,t) dx = 1$.

We've done this in class too!

Section 2.4 Q18: Solve the heat eq with convection:

$$\begin{cases} u_t - ku_{xx} + Vu_x = 0 & t > 0 \quad -\infty < x < \infty \\ u(x,0) = \phi(x) & -\infty < x < \infty \end{cases}$$

where V is a constant.

Make the substitution $y = x - Vt$, $x = y + Vt$:

Define $v(y,t) = u(y+Vt, t)$.

Then $v_t = u_x \cdot V + u_t$

$$v_x = u_x$$

$$v_{xx} = u_{xx}$$

So: $0 = \underbrace{u_t + Vu_x}_{v_t} - k \underbrace{u_{xx}}_{v_{xx}} = v_t - kv_{xx}$

So v satisfies the usual diffusion eq. with the initial condition $v(y,0) = u(y,0) = \phi(y)$. Hence

$$v(y,t) = \int_{-\infty}^{\infty} \zeta(y-w, t) \phi(w) dw$$

$$\rightarrow u(x,t) = v(x-Vt, t) = \int_{-\infty}^{\infty} \zeta(x-Vt-w, t) \phi(w) dw$$