## 5.2 Algebra of limits

Given some functions  $f : \mathbb{R} \to \mathbb{R}$  and  $g : \mathbb{R} \to \mathbb{R}$ , our goal is to understand how limits of  $f \pm g$  might behave, as well as limits of  $f \cdot g$  and  $\frac{f}{g}$ . In general we should exercise caution!

**Example 5.1:** Let  $C \in \mathbb{R}$ . Consider the sequences  $a_n = n + C$  and  $b_n = -n$ . Then:

$$\lim_{n\to\infty} a_n = +\infty \qquad \lim_{n\to\infty} b_n = -\infty$$

and

$$\lim_{n\to\infty} (a_n + b_n) = \lim_{n\to\infty} (n + C - n) = \lim_{n\to\infty} C = C.$$

This suggests that  $+\infty - \infty = C$  for any C. The analogous example for functions would involve f(x) = x + C and g(x) = -x.

**Example 5.2:** Consider the sequences  $a_n = 2n$  and  $b_n = -n$ . Then:

$$\lim_{n\to\infty} a_n = +\infty \qquad \lim_{n\to\infty} b_n = -\infty$$

and

$$\lim_{n\to\infty}(a_n+b_n)=\lim_{n\to\infty}(2n-n)=\lim_{n\to\infty}n=+\infty.$$

**Example 5.3:** Consider the sequences  $a_n = n$  and  $b_n = -2n$ . Then:

$$\lim_{n\to\infty}a_n=+\infty\qquad \lim_{n\to\infty}b_n=-\infty$$

and

$$\lim_{n\to\infty}(a_n+b_n)=\lim_{n\to\infty}(n-2n)=\lim_{n\to\infty}(-n)=-\infty.$$

These examples demonstrate that we should be cautious when taking limits of sums or differences of sequences and functions. In particular, these examples show that  $+\infty-\infty$  is a meaningless expression. Similar absurdities can be achieved by multiplication or division of sequences or functions. Let's list the meaningless expressions:

## Meaningless expressions

The following expressions do not make sense:

$$+\infty - \infty$$
  $-\infty + \infty$   $\pm \infty \cdot 0$   $\frac{\pm \infty}{\pm \infty}$   $\frac{0}{0}$ 

In contrast, here are the meaningful expressions:

### Meaningful expressions

The following expressions do make sense:

$$+\infty + C = +\infty \quad (\text{if } C \in \mathbb{R} \cup \{+\infty\})$$

$$-\infty + C = -\infty \quad (\text{if } C \in \{-\infty\} \cup \mathbb{R})$$

$$\pm \infty \cdot C = \pm \infty \quad (\text{if } C \in \mathbb{R}_+ \cup \{+\infty\})$$

$$\pm \infty \cdot C = \mp \infty \quad (\text{if } C \in \{-\infty\} \cup \mathbb{R}_-)$$

$$\frac{\pm \infty}{C} = \pm \infty \quad (\text{if } C \in \mathbb{R}_+)$$

$$\frac{\pm \infty}{C} = \mp \infty \quad (\text{if } C \in \mathbb{R}_-)$$

$$\frac{C}{\pm \infty} = 0 \quad (\text{if } C \in \mathbb{R})$$

A more delicate meaningful expression is

$$\frac{C}{0} = \infty \quad (\text{if } C \in \{-\infty\} \cup \mathbb{R} \cup \{+\infty\} \setminus \{0\})$$

However to determine whether it is  $+\infty$  or  $-\infty$  we need to look at the numerator and the denominator in a neighborhood of the point in question. This could lead to different left and right limits (in the case that the denominator changes sign at the point in question).

#### **Example 5.4:** We start with a simple example:

$$f(x)=\frac{1}{x}.$$

At x = 0, we have an expression of the form  $\frac{C}{0}$ . The constant C = 1 > 0 in this case, and the denominator changes sign from negative (left of 0) to positive (right of 0). Hence

$$\lim_{x \to 0^{-}} \frac{1}{x} = -\infty \quad \text{and} \quad \lim_{x \to 0^{+}} \frac{1}{x} = +\infty.$$

#### **Example 5.5:** We analyze the rational function

$$f(x) = \frac{x - 2}{x^2 - 2x + 1}.$$

Numerator: x - 2 is < 0 when x < 2, = 0 when x = 2 and > 0 when x > 2. Denominator:  $x^2 - 2x + 1 = (x - 1)^2$  is always  $\ge 0$ , and = 0 only at x = 1.

At x = 1 we encounter the situation  $\frac{C}{0}$  with C < 0. Near x = 1, the denominator is > 0, so that both left and right limits must be  $-\infty$  and we have:

$$\lim_{x \to 1} \frac{x - 2}{x^2 - 2x + 1} = -\infty.$$

The rules we stated allow us to state the following important theorem:

**Theorem 5.5** (Algebra of Limits): Let  $f : \mathbb{R} \to \mathbb{R}$  and  $g : \mathbb{R} \to \mathbb{R}$ , and let  $x_0$  either be a real number, or  $+\infty$  or  $-\infty$ . Suppose that as  $x \to x_0$ , both f and g have limits

$$\lim_{x \to x_0} f(x) = \ell_f \quad \text{and} \quad \lim_{x \to x_0} f(x) = \ell_g$$

(these limits can be finite of infinite). Then the following equalities hold *only* when the right hand side is a meaningful expression:

$$\lim_{x \to x_0} (f(x) \pm g(x)) = \ell_f \pm \ell_g$$

$$\lim_{x \to x_0} (f(x) \cdot g(x)) = \ell_f \cdot \ell_g$$

$$\lim_{x \to x_0} \frac{f(x)}{g(x)} = \frac{\ell_f}{\ell_g},$$

where in the last case, g(x) must be nonzero for all x tending to  $x_0$ , and it is possible that there are different left and right limits (for instance, if g changes sign at  $x_0$ ).

*Proof.* We prove just one case to demonstrate the idea of the proof. Let us prove that

$$\lim_{x \to x_0} (f(x) + g(x)) = \ell_f + \ell_g$$

in the case that both  $\ell_f$  and  $\ell_g$  are real numbers (and not  $\pm \infty$ ). That is, we need to show that for any  $\varepsilon > 0$  there exists  $\delta = \delta(\varepsilon) > 0$  such that for all  $0 < |x - x_0| < \delta$ , we have that  $|(f(x) + g(x)) - (\ell_f + \ell_g)| < \varepsilon$ .

Fix  $\tilde{\varepsilon} > 0$ . We start with f: since  $\lim_{x \to x_0} f(x) = \ell_f$ ,

$$\exists \delta_f = \delta_f(\tilde{\varepsilon}) > 0 \text{ such that } \forall 0 < |x - x_0| < \delta_f, |f(x) - \ell_f| < \tilde{\varepsilon}.$$

Similarly, for g, using the same  $\tilde{\epsilon}$ :

$$\exists \delta_g = \delta_g(\tilde{\epsilon}) > 0 \text{ such that } \forall 0 < |x - x_0| < \delta_g, |g(x) - \ell_g| < \tilde{\epsilon}.$$

Define  $\delta = \min\{\delta_f, \delta_g\}$ . Then for all  $0 < |x - x_0| < \delta$ ,

$$\begin{aligned} |(f(x)+g(x))-(\ell_f+\ell_g)| &= |(f(x)-\ell_f)+(g(x)-\ell_g)|\\ (triangle\ inequality) &\leq |f(x)-\ell_f|+|g(x)-\ell_g|\\ &< \tilde{\varepsilon}+\tilde{\varepsilon}=2\tilde{\varepsilon}. \end{aligned}$$

By defining  $\varepsilon = 2\tilde{\varepsilon}$  we are done: we have shown that for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that for all  $0 < |x - x_0| < \delta$ , we have  $|\underbrace{(f(x) + g(x))}_{\text{this is the function}} - \underbrace{(\ell_f + \ell_g)}_{\text{this is the limit}} | < \varepsilon.$ 

An immediate corollary is this:

**Corollary 5.6:** If f and g are continuous at  $x_0 \in \mathbb{R}$ , then so are  $f \pm g$ ,  $f \cdot g$  and  $\frac{f}{g}$  continuous at  $x_0$  (the last one only if  $g(x_0) \neq 0$ ).

Another corollary is this:

**Corollary 5.7:** Rational functions are continuous on their domains and polynomials are continuous on  $\mathbb{R}$ .

# 5.3 Comparison theorems

**Theorem 5.8:** Let  $f : \mathbb{R} \to \mathbb{R}$  and  $g : \mathbb{R} \to \mathbb{R}$  and let  $x_0 \in \{-\infty\} \cup \mathbb{R} \cup \{+\infty\}$ . Suppose that

$$\lim_{x \to x_0} f(x) = \ell_f \quad \text{and} \quad \lim_{x \to x_0} g(x) = \ell_g.$$

If in a neighborhood of  $x_0$  (excluding  $x_0$  itself) we have  $f(x) \le g(x)$ , then  $\ell_f \le \ell_g$ .

*Proof.* We skip the proof here. It can be found in the book.

**Theorem 5.9** (Squeeze Theorem): Let  $f, g, h : \mathbb{R} \to \mathbb{R}$  and let  $x_0 \in \{-\infty\} \cup \mathbb{R} \cup \{+\infty\}$ . If

$$\lim_{x \to x_0} f(x) = \lim_{x \to x_0} h(x) = \ell \in \mathbb{R} \quad \text{and} \quad f \le g \le h \text{ in a neighborhood of } x_0$$
(excluding  $x_0$  itself)


then

$$\lim_{x \to x_0} g(x) = \ell.$$

*Proof.* We skip the proof here. It can be found in the book.

**Example 5.6:** Using this theorem we can immediately conclude that

$$\lim_{x \to 0} x^2 \sin\left(\frac{1}{x}\right) = 0.$$



**Example 5.7:** The Squeeze Theorem justifies our computation in Example 4.12 (the removable singularity of  $\frac{\sin x}{x}$  at x = 0). In that problem we found that

$$\underbrace{\cos x}_{f(x)} < \underbrace{\frac{\sin x}{x}}_{o(x)} < \underbrace{\frac{1}{h(x)}}$$

and used the fact that  $\lim_{x\to 0} \cos x = 1$  to conclude that

$$\lim_{x \to 0} \frac{\sin x}{x} = 1.$$